

CONSTRUCTION OF ADCRE VECTORS




FIGURE 1

EXPRESSION OF CRE RECOMBINASE IN CELLS INFECTED WITH ADCRE





FIGURE 2

CONSTRUCTION OF AD VECTORS EXPRESSING LUCIFERASE UNDER CONTROL OF A MOLECULAR SWITCH

A



B





FIGURE 3

CRE MEDIATED RECOMBINATION

A



B





Figure 4

EXCISION OF SEQUENCES FROM ADMA19 IN CELLS COINFECTED WITH ADMA19 AND ADCRE

A



B





FIGURE 5A,B

EXPRESSION OF LUCIFERASE IN AD VECTOR INFECTED CELLS

Expt. 1



Expt. 2





FIGURE 5C


STRUCTURE OF INTEGRATED SEQUENCES IN CELL LINE PW27C2



EXPRESSION OF NEO RESULTING IN G418 RESISTANCE

FIGURE 6A

STRUCTURE OF INTEGRATED SEQUENCES IN CELL LINE PW27C2



EXPRESSION OF NEO RESULTING IN G418 RESISTANCE

FIGURE 6A

SOUTHERN BLOT HYBRIDIZATION ANALYSIS OF DNA FROM CELL LINES INFECTED WITH ADCRE




FIGURE 6C

CRE-MEDIATED INSERTION OF A PLASMID ENCODING HISD SEQUENCES INTO THE LOX SITE OF CELL LINE PW27C2




FIGURE 7A

**SOUTHERN BLOT HYBRIDIZATION ANALYSIS OF CELL LINES
DERIVED BY CRE MEDIATED INTEGRATION OF pPW1**




FIGURE 7B

**SOUTHERN BLOT HYBRIDIZATION ANALYSIS OF CELL LINES
DERIVED BY CRE MEDIATED INTEGRATION OF pPW1**





FIGURE 7B

CRE-MEDIATED EXCISION OF DNA CONTAINING hisD SEQUENCES FLANKED BY LOX SITES



**Loss of hisD expression (Histidinol sensitive)
Gain of neo expression (G418 resistant)**

FIGURE 8A

Conversion of HisD resistant cells to
G418 resistant cells by infection with AdCre






FIGURE 8B

IN VIVO GENE EXPRESSION CONTROLLED BY A MOLECULAR SWITCH



TRANSGENICS CONTAINING GENES CONTROLLED BY A MOLECULAR SWITCH



EXPRESSION OF β -Gal, Rb, P53, Neu ETC.

Figure 9